

Subject ID:	Rater:	Date:
Age at	Interview:	Apparent onset:
		First treatment:

Duration:	0 - Absent	1 - < 2 wk (≥ hrs)	2 - 2+ wk	3 - 2+ mo or 2+ ep of ≥ 2 wk	4 - 2+ years
Severity:	0 - Absent	1 - Minimal. Very mild sx or only suspected	2 - Moderate. Definite, clinically significant sx.	3 - Severe. Clearly interferes with function or preoccupies	4 - Very severe. Gross or nearly constant effect on function

DURATION SEVERITY

- | | | |
|-----------|-----------|--|
| 0 1 2 3 4 | 0 1 2 3 4 | P-1. Any delusions. Any type, not limited to the following (check all that apply):
___a. Paranoid ___b. Non-paranoid ___c. Grandiose ___d. Somatic ___e. Religious ___f. Nihilistic
___g. Delusions accompanied by hallucinations for at least one week |
| 0 1 2 3 4 | 0 1 2 3 4 | P-2. Paranoia (preoccupation with persecutory ideas, delusional or not) |
| 0 1 2 3 4 | 0 1 2 3 4 | P-3. Any hallucinations. (Abnormal full sensory perception when awake.) Any type, not limited to the following:
___a. Non-affective verbal hallucinations spoken to the subject |
| 0 1 2 3 4 | 0 1 2 3 4 | S-1. Control delusions. ___a. Thought insertion ___b. Thought withdrawal ___c. Control of thought/actions |
| 0 1 2 3 4 | 0 1 2 3 4 | S-2. Other bizarre delusions. (Implausible, impossible, other than control delusions) |
| 0 1 2 3 4 | 0 1 2 3 4 | S-3. Conversing/commenting/continuous hallucinations.
___a. Voices conversing ___b. Voices commenting ___c. Continuous (throughout the day for several days) |
| 0 1 2 3 4 | 0 1 2 3 4 | S-4. Abnormal perception of thought.
___a. Thought broadcasting ___b. Audible thoughts ___c. Thought echo |
| 0 1 2 3 4 | 0 1 2 3 4 | NAP. Psychosis (2+ weeks) without prominent mood symptoms:
___a. After mania ___b. After depression ___c. Neither |
| 0 1 2 3 4 | 0 1 2 3 4 | N-1. Blunted (restricted) affect. |
| 0 1 2 3 4 | 0 1 2 3 4 | N-2. Poverty of speech. (Negative thought disorder) |
| 0 1 2 3 4 | 0 1 2 3 4 | D-1. Formal thought disorder. (Impaired understandability; abnormal use of words) |
| 0 1 2 3 4 | 0 1 2 3 4 | D-2. Bizarre behavior. (Grossly unusual dress or social/sexual/agitated/ritualistic behavior likely due to psychosis) |
| 0 1 2 3 4 | 0 1 2 3 4 | DE-1. Depression. (Dysphoria and associated symptoms. Severity of 3-4 requires full major depressive syndrome.) |
| 0 1 2 3 4 | 0 1 2 3 4 | DE-2. Maximum # of 9 depressive features (a-i below) concurrent for 2+ weeks ever. |
| 0 1 2 3 4 | 0 1 2 3 4 | M-1. Mania. (Manic mood and associated symptoms. Severity of 3-4 requires full manic syndrome.) |
| 0 1 2 3 4 | 0 1 2 3 4 | M-2. Maximum # of 7 manic features other than mood (a-i below) concurrent for 1+ weeks ever (-1 if only irritable) |
| 0 1 2 3 4 | 0 1 2 3 4 | MP-1. Concurrent depressed mood + delusions or hallucinations of:
___a. Guilt ___b. Catastrophe/nihilism ___c. Suicide ___d. Disease (non-bizarre) |
| 0 1 2 3 4 | 0 1 2 3 4 | MP-2. Concurrent grandiose/manic mood + delusions or hallucinations of:
___a. Importance/power ___b. Special relationship with God /mission |
| 0 1 2 3 4 | 0 1 2 3 4 | C. Deterioration. Residuum between exacerbations is: 0-Absent 1-Mild 2-Significant 3-Severe 4-Severe, > 5 yr |
| 0 1 2 | 0 1 2 | A. Complicating factors are 0-Absent 1-Present but unlikely to be causative 2-Strongly suspected or likely causes of psychosis. Specify: _____
___a. Substance abuse ___b. Organic ___c. Atypical features ___d. Personality disorder ___e. Dissociative |
| 1 2 3 4 | | Quality of available information.
4 -Excellent 3 - Minor inadequacies 2 - Major inadequacies/reduced confidence 1 - Very limited |

Depression and mania checklists are provided to assist in reviewing cases, Use ticks to not presence in different episodes. Use next line to note rated episodes.

___ a. Depressed mood. ___ b. Markedly diminished pleasure in most/all activities. ___ c. Increased or decreased weight (>5%) or appetite. ___ d. Increased/decreased sleep. ___ e. Observable psychomotor agitation or retardation. ___ f. Fatigue or loss of energy. ___ g. Worthlessness or excessive/inappropriate guilt (not just about being sick). ___ h. Diminished ability to think/concentrate/ indecisiveness , due to depression. ___ i. Recurrent thoughts of death or suicide , or suicide attempt or plan.	___ x. Abnormally and persistently elevated or expansive mood. ___ y. Abnormally and persistently irritable mood (with excitement or agitation). ___ a. Inflated self-esteem or grandiosity. ___ b. Decreased need for sleep. ___ c. More talkative than usual or pressure to keep talking. ___ d. Flight of ideas or subjectively racing thoughts. ___ e. Distractibility (attention too easily drawn to unimportant/irrelevant external stimuli). ___ f. Increased goal-directed activity (social/sexual/work) or psychomotor agitation. ___ g. Excessive pleasurable activities with high potential for painful consequences.
--	---